












The Physicochemical Analysis of the Interaction Reaction of Potassium Chloride with Ammonium Hydrosulfate



Kalkaman Zhumashev¹ , Akmaral Serikbayeva^{2,*} , Akkenzhe Bussurmanova² , Anar Akkenzheyeva² , Assiya Boranbayeva² , Zhansaule Altybayeva² , Aigul Gusmanova² , Aibala Narembekova³ , Feruza Berdikulova⁴ , Arman Mauleshev⁵  and Tugelbai Kenbaev⁵ 

¹«Innovation» LLP, Karaganda, Kazakhstan

²Caspian University of Technology and Engineering, named after Sh. Yessenov, Aktau, Kazakhstan

³Karaganda Technical University named after A. Saginov, Karaganda, Kazakhstan

⁴RSE «National Center for Complex Processing of Mineral Raw Materials of the Republic of Kazakhstan,» Almaty, Kazakhstan

⁵«KazAzot» JSC, Aktau, Kazakhstan

Abstract:

Introduction: This paper represents the results of exploratory research on the production of ammonium-potassium sulfate salt and by-products. Information on the use of the results of fundamental research for the creation of new technological processes is provided. The increase in the range of products based on ammonium salts is a priority task for the industrial enterprise "KazAzot" JSC (Kazakhstan, Aktau). Another task is to expand the areas of application of products with derivatives based on ammonium hydrosulfate. Furthermore, combined fertilizers can be included in the list of products. The aim of the work is to scientifically substantiate physicochemical transformations in reaction mixtures, such as obtaining ammonium-potassium sulfate-fertilizer with three target components.

Methods: The experimental work was conducted in a laboratory shaft electric furnace, with observations of changes in the reaction mass and determination of the pH of the gas phase. The produced products were subjected to an X-ray phase analysis on the diffractometer Miflex 600 (Japan) with CuK α . In addition, the physicochemical changes of reaction mixtures and synthesized salts were studied with the help of the derivatograph of the Q-1000/D system by "MOM" F. Paulik, J. Paulik, L. Erdey and LabSys Evo TG-DTA/DTG (Setaram, France), NETZSCH STA 443 F3 in the nitrogen atmosphere. The thermodynamic evaluation of the probability of reactions was obtained using the Astra-4 program complex HCS-5.1 Chemistry (Outokumpu).

Results: It has been established that in the temperature range 157 - 260 °C at equal molar ratios of NH₄HSO₄ ÷ KSO₄ the condensed product of the reaction is a double salt - ammonium-potassium sulfate (NH₄KSO₄) and its decomposition into constituent sulfates occurs above 260 °C with simultaneous pyrolysis of ammonium sulfate to complete formation of ammonium hydrosulfate and ammonia. At incomplete pyrolysis of ammonium sulfate in the cooled residue is present (NH₄)₃HS₂O₈, so salt formed from 2 ammonium sulfates: (NH₄)₂SO₄•NH₄HSO₄ and similarly, the product of incomplete decomposition of ammonium-potassium sulfate - NH₄KSO₄ is KHSO₄•K₂SO₄.

Conclusion: An increase in the stoichiometric molar ratio in the system «NH₄HSO₄ ÷ KCl» leads to the formation of more complex hydrated salts and the basic salt of ammonium - ammonium sulfate, and, on the contrary, a decrease in the ratio - to a more basic compound of potassium sulfate.

Keywords: Potassium chloride, Thermodynamic analysis, Reaction mechanism, Hydrosulfates, Ammonia, Ammonium sulfate.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Received: May 10, 2024

Revised: July 18, 2024

Accepted: August 23, 2024

Published: September 12, 2024



Send Orders for Reprints to
reprints@benthamscience.net

*Address correspondence to this author at the Caspian University of Technology and Engineering, named after Sh. Yessenov, Aktau, Kazakhstan; E-mail: akmaral.serikbayeva@yu.edu.kz

Cite as: Zhumashev K, Serikbayeva A, Bussurmanova A, Akkenzheyeva A, Boranbayeva A, Altybayeva Z, Gusmanova A, Narembekova A, Berdikulova F, Mauleshev A, Kenbaev T. The Physicochemical Analysis of the Interaction Reaction of Potassium Chloride with Ammonium Hydrosulfate. *Open Chem Eng J*, 2024; 18: e18741231321361.

<http://dx.doi.org/10.2174/0118741231321361240905053845>

1. INTRODUCTION

Ammonium sulfate is widely used in various spheres, mainly in agriculture as a fertilizer, as it is a source of nitrogen and sulfur. It contains up to 21% nitrogen, which has a beneficial effect on plants, primarily on the root system of plants [1-3].

Compared to other nitrogen fertilizers such as urea and ammonium nitrate, ammonium sulfate has some agronomic and environmental advantages [4-6]. Researchers [7, 8] have developed a method for obtaining ammonium sulfate in the form of granules, and it is used both individually and as part of compound fertilizers. Obtaining ammonium sulfate by this method decreases the size of granules and increases their strength.

However, despite the known recommendations on the use of ammonium sulfate for growing crops (from potatoes to citrus) on chernozem and sulfur soils, it is used only to a limited extent. The reason for this is the availability of complex fertilizers richer in nitrogen and other useful components. Another reason is the acidic property of fertilizers, which can lead to rapid acidification of the soil. However, the lands of the Mangistau region contain limestone - an alkaline reagent and therefore, the use of fertilizer based on ammonium sulfate will not have a harmful effect on the soil [9].

Unfortunately, we did not find any work on the study of the solid-phase interaction of ammonium sulfate with potassium chloride during the search. All studies on the production of potassium sulfate from various mixtures, such as sodium sulfate and potassium chloride, are devoted to aqueous solutions [10-12]. As an analogue, the interaction between ammonium bisulfate and ammonium chloride can be cited. The reaction products are hydrogen chloride and ammonium sulfate at a temperature below the initial pyrolysis point of 260 °C of the latter compound. The results of the study do not take into account the physicochemical transformations in reaction mixtures before the formation of the final thermally stable compound, potassium sulfate.

Furthermore, it is relevant to obtain double salts on the basis of ammonium sulfate, which is easily assimilated by plants [13].

Ammonium salt-based product range increase is a priority task for the industrial enterprise JSC «KazAzot» (Kazakhstan, Aktau). In order to solve this task, the

technology of granulated ammonium sulfate production is being mastered by the enterprise. Despite the fact that the methods of obtaining granules of these fertilizers are known [14-16], the management of the enterprise sets new tasks for researchers - to search for the possibility of using binders based on their own products.

The other objective is to expand the areas of application of products with derivatives based on ammonium sulfate [17-19]. These include combined fertilizers based on ammonium hydrosulfate. The authors have previously studied and given the results of physicochemical analysis of a number of topochemical reactions of the interaction of metal compounds with ammonium sulfate [20].

An analysis of the scientific and patent literature [21, 22] represented that to obtain complex mineral fertilizers as a nitrogen-containing component mainly use ammonium nitrate and urea, which have a high cost. There is limited information in scientific and patent literature on the use of ammonium sulfate for the production of complex fertilizers.

The research aims to provide a scientific basis for the possibility of achieving the set objectives by studying the physicochemical transformations in reaction mixtures for example, to obtain ammonium-potassium sulfate - a fertilizer with three target components.

2. MATERIALS AND METHODS

The experimental research was carried out in a laboratory shaft electric furnace to study the interaction of potassium chloride with ammonium hydrosulfate with observations of changes in the reaction mass and determination of the pH of the gas phase. It was a DTA installation with a transformer and a voltage regulator at a speed of 10 °/minute. The samples were placed in an oven in a crucible made of heat-resistant glass. Additionally, when the maximum temperature is set, the heating is turned off using a thermostat equipped with a chromel-alumel thermocouple.

The samples were placed in the furnace in a crucible made of heat-resistant glass. The required temperature in the furnace was kept constant using a device equipped with a chromel-alumel thermocouple. In all experiments, the sample weight of the substances and reaction mixtures used for the study was calculated according to the assumed reaction equations. Depending on the purpose of

the experiment, the sample holding time in the oven was within 60 minutes. The ammonium hydrosulfate of the "clean" label was used for the experiment.

The experimental products were subjected to an X-ray phase analysis on a Miflex 600 diffractometer (Japan) with $\text{CuK}\alpha$ at a scanning range of 3° to 90° , and a step of 0.02. Physicochemical changes of the reaction mixtures and some synthesized salts upon heating were studied using a Q-1000/D derivatograph system Q-1000/D by "MOM" F. Paulik, J. Paulik, L. Erdey and LabSys Evo TG-DTA/DTG (Setaram, France), NETZSCH STA 443 F3 in nitrogen atmosphere at a feed rate of 50ml/minute at a heating rate of 10° per minute. The thermodynamic evaluation of reaction probability was obtained using the HCS-5.1 Chemistry (Outokumpu) "Astra-4" software package HCS-5.1 Chemistry (Outokumpu) [23] (Table 1). The appropriate thermodynamic functions of hydrosulfates for

which no thermodynamic data was available were calculated by the ionic increment method [24] in order to estimate the probability of their formation.

The comparison of reference data of a number of known compounds, such as ammonium sulfate, with the calculated values of thermodynamic values by the method of ionic increments showed no more than a 1% discrepancy (Table 2).

The charge was made according to the reaction equation with an excess of ammonium hydrosulfate 18.9738g (10.0%), a total of 134.084g, and the mass of potassium chloride in the charge was equal to 74.615g, grade "clean for analysis" with the content of the main substance 99.98%. The total calculated mass of the charge according to the equation of reaction with excess ammonium hydrosulfate was 208.699g. The samples for research were taken from the following mixture.

Table 1. Thermodynamic functions of complex ammonium salts calculated using reference values of ionic increments at 298K.

Compounds	ΔH , kCal	ΔG , kCal	ΔS , kCal
$(\text{NH}_4)_2\text{SO}_4(\text{cr.})$	-1180,0	-901,3	220
NH_4HSO_4	-1020,2	-834,7	242,4
$(\text{NH}_4)_3\text{H}(\text{SO}_4)_2$	- 2215,8	- 1726,5	361,4
$\text{NH}_4\text{K}(\text{HSO}_4)_2$	- 2159,2	- 1871,2	472,9
$\text{NH}_4\text{K}_2(\text{HSO}_4)_3$	- 3298,2	- 2907,7	703,4
$\text{NH}_4\text{K}_2\text{HS}_2\text{O}_8$	- 2431,9	- 2141,3	465,6
$(\text{NH}_4)_2\text{KHS}_2\text{O}_8$	- 2313,3	- 1939,5	477,5
KHSO_4	- 1139,0	- 1036,5	230,5
NH_4KSO_4	- 1292,9	- 1104,8	235,1
$\text{K}_3\text{H}(\text{SO}_4)_2$	- 2572,2	- 2331,9	343,9
$\text{K}_3\text{NH}_4(\text{SO}_4)_2$	- 2704,6	- 2411,4	458,3
$\text{K}_2\text{CO}_3(\text{cr.})$	-1146,1	-1059,8	156,3
$\text{K}_2\text{SO}_4(\text{cr.})$	-1433,7	-1316,4	175,7
$\text{K}_2\text{S}_2\text{O}_7(\text{cr.})$	- 2069,3	-1904,4	340,5

Table 2. The results of values of thermodynamic functions of the assumed variants of chemical interaction of potassium salts with ammonium hydrosulfate.

No.	An Equation of a Chemical Reaction	$T_{\text{pr}}^\circ\text{C}$	ΔH_{298}	ΔG_{298}	ΔS_{298}
1	$\text{NH}_4\text{HSO}_4 + \text{KCl} = \text{HCl} + \text{NH}_4\text{KSO}_4$	205,5	71,4	43,1	96,9
2	$\text{NH}_4\text{KSO}_4 = \frac{1}{2} (\text{K}_2\text{SO}_4 + (\text{NH}_4)_2\text{SO}_4)$	101,5	- 13,95	- 4,05	- 37,25
3	$\frac{1}{2} (\text{NH}_4)_2\text{SO}_4 = \frac{1}{2} (\text{NH}_4\text{HSO}_4 + \text{NH}_3)$	255	56,8	24,95	107,5
4	$\frac{1}{2} (\text{K}_2\text{SO}_4 + \text{NH}_4\text{HSO}_4) = \frac{1}{2} (\text{K}_2\text{S}_2\text{O}_7 + \text{H}_2\text{O} + \text{NH}_3)$	27,5	178,95	109	118,8
Σ_{1-4}	$\text{NH}_4\text{HSO}_4 + \text{KCl} = \text{HCl} + \frac{1}{2} (\text{K}_2\text{S}_2\text{O}_7 + \text{H}_2\text{O}) + \text{NH}_3$	481	293,2	176	388,75
Possible reactions of formation of acid salts					
5	$\text{K}_2\text{SO}_4 + 2\text{NH}_4\text{HSO}_4 = 2\text{KHSO}_4 + (\text{NH}_4)_2\text{SO}_4$	- 3	- 11,2	0,7	- 41,5
6	$3\text{NH}_4\text{KSO}_4 + \text{NH}_4\text{HSO}_4 = (\text{NH}_4)_3\text{HS}_2\text{O}_8 + \text{K}_3(\text{HS}_2\text{O}_8) + \text{NH}_3$	-78	64,7	74	332,2
7	$3\text{K}_2\text{SO}_4 + 2\text{NH}_4\text{HSO}_4 = 2\text{K}_3(\text{HS}_2\text{O}_8) + (\text{NH}_4)_2\text{SO}_4$	- 50	- 64,8	21,1	-290,1
8	$(\text{NH}_4)_2\text{SO}_4 + \text{NH}_4\text{HSO}_4 = (\text{NH}_4)_3\text{HS}_2\text{O}_8$	- 118	- 15,6	9,5	- 101
9	$3\text{NH}_4\text{HSO}_4 + \text{NH}_4\text{K}_2\text{HS}_2\text{O}_8 = 2\text{NH}_4\text{K}(\text{HSO}_4)_2 + (\text{NH}_4)_2\text{SO}_4$	-54,5	- 5,9	1,7	- 27

3. RESULTS AND DISCUSSION

The article presents the research data aimed at creating a technological basis for the processing of metallurgical raw materials using ammonium hydrosulfate. However, it is possible that these reactions are summarized for successive physicochemical transformations with intermediate products stable up to a certain temperature. Such compounds can include ammonium-metal sulfates - anhydrous salts of Tutton, and hydrosulfates. The calculated values of thermodynamic functions of some complex ammonium salts at 298K are presented in Table 1.

The results of the values of thermodynamic functions of the assumed variants of chemical interaction of potassium salts with ammonium hydrosulfate melt are presented in Table 2.

Since weight loss was calculated as a percentage of potassium chloride, the calculation was carried out according to the reaction equation 1 with respect to the charge, weighing 189.725g. The excess of ammonium hydrosulfate at the temperatures of its decomposition was taken into account separately. The comparative values of calculation by the equations of assumed reactions and data on TG are given in Table 3 and Fig.(1).

Table 3. Mass loss results are calculated from reactions and TG curve readings.

N ^o	Reaction Equations	t ₁ - t ₂ , °C	Δm _{calculation} %	Δm _{fact} %
0	NH ₄ HSO ₄ (s) = NH ₄ HSO ₄ (l)	30 - 157.3	0	0
1	NH ₄ HSO ₄ + KCl = HCl + NH ₄ KSO ₄	157.3-381	19.254	20.26-1.006 = 19.254
2	NH ₄ KSO ₄ = ½ (K ₂ SO ₄ + (NH ₄) ₂ SO ₄)	381- 381		
3	½ (NH ₄) ₂ SO ₄ = ½ (NH ₄ HSO ₄ (l) + NH ₃) [8]	260 - 360	4.4838	4.4838
4	½ (K ₂ SO ₄ + NH ₄ HSO ₄) = ½ (K ₂ S ₂ O ₇ + H ₂ O + NH ₃)	381 - 450	9.2314	32.5-23.7378 = 8.7622
Σ	NH ₄ HSO ₄ + KCl = ½ (K ₂ S ₂ O ₇ + H ₂ O) + NH ₃ + HCl	30 - 450	32.9692	32.5
9	3NH ₄ HSO ₄ + NH ₄ K ₂ HS ₂ O ₈ = 2NH ₄ K(HSO ₄) ₂ + (NH ₄) ₂ SO ₄	260 - 190	8.0	8.95
	NH ₄ HSO ₄ = SO ₃ + NH ₃ + H ₂ O [8]	450 - 380	9.38	41.88 - 32.5 = 9.38




Fig. (1). DTA and TG curves of interaction between potassium chloride and ammonium hydrosulfate.

In the temperature range of 125 - 157 °C, the DTA curve has a rectilinear section, characteristic of the melting of ammonium hydrosulfate. In this section, according to TG readings, there is no gas emission.

In the interval of temperatures 157 - 381 °C on the DTA curve, a strong endothermic effect is fixed, which corresponds to the loss of mass by TG 20.26% of the mass of the charge suspension. Furthermore, the calculation according to the equation of reaction 1, the condensed product of the reaction is a double salt - ammonium-potassium sulfate - NH_4KSO_4 , and the mass of released hydrogen chloride by potassium chloride in the charge should correspond to 19.254%. When it decomposes by reaction 2, there should be no loss of mass. Therefore, the difference of 1.006% can be attributed to the beginning of ammonia release as a result of the primary pyrolysis of ammonium sulfate by reaction 3. It should be noted that, in fact, reaction 1 is the initial and limiting reaction for successive reactions 2 and 3, despite the greater probability of the latter occurring. Therefore, as NH_4KSO_4 appears, its decomposition into constituent sulfates occurs, and in the temperature range of 235 - 340 °C, the pyrolysis of ammonium sulfate to the formation of ammonium hydrosulfate and ammonia is completed [25, 26]. It is likely that in the studied mixture, pyrolysis starts earlier due to the release of heat energy via reaction 2.

The amount of gases released as the maximum heating temperature of - 450 °C reached according to the TG curve was $32.5 - 20.26 + 1.006 = 13.246\%$. Reaction 4 failed to complete at 5.083%.

The additional outgassing occurs at the cooling stage due to the completion of reaction 4 and depends on the stoichiometric ratios of the components of the direct reaction products according to one of the reaction equations 5 - 9. At incomplete pyrolysis of ammonium sulfate, XRA of the cooled residue shows the presence of $(\text{NH}_4)_3\text{HS}_2\text{O}_8$, so the salt formed from 2 ammonium sulfates: $(\text{NH}_4)_2\text{SO}_4 \cdot \text{NH}_4\text{HSO}_4$ [12]. Depending on the sensitivity of the device and the weight of the sample, the beginning of pyrolysis on the DTA curves is fixed in the range of 230-260 °C.

The precipitation of such salts from solution is generally known. Similarly, the incomplete decomposition product of ammonium-potassium sulfate - NH_4KSO_4 is $\text{KHSO}_4 \cdot \text{K}_2\text{SO}_4$. These salts consist of a basic salt and an acidic salt. Therefore, their formulas should be written as compounds containing an anion such as hydrosulfate and sulfate: $(\text{NH}_4)_3\text{HS}_2\text{O}_8$ and $\text{K}_3(\text{HS}_2\text{O}_8)$. As shown by the DTA data (Table 2 and Fig. 1) of the cooled residues, they

contain a compound, double hydrosulfate- $2\text{NH}_4\text{K}(\text{HSO}_4)_2$, which was formed by reaction 7.

An excess of potassium pyrosulfate of reaction 5 is also retained during cooling. Therefore, because of the lack of ammonium hydrosulfate, the reaction starts with the formation of normal double salt - ammonium potassium sulfate, which, in the course of temperature increase, undergoes successive physicochemical transformations. In this case, at certain temperature intervals, the products can be stable compounds, as well as the probable short-term appearance of thermally unstable acidic salts. The formation of acid salts at the heating stage of potassium salts and ammonium hydrosulfate is the result of the decomposition of double sulfate and their intermediate products, which are accompanied by the formation of ammonium sulfate or hydrosulfates on its basis. Above the pyrolysis temperature of ammonium sulfate, the reactions proceed with the liberation of ammonia and the decomposition of the double acid salts into ammonium hydrosulfate and potassium sulfate. When such a condensed mixture of direct reactions is cooled below 260 °C, acidic salts and ammonium sulfate or double acidic salts are formed again, depending on the stoichiometry of the initial mixtures. The process can continue with the release of ammonia until this temperature is reached.

When the stoichiometric molar ratio of $\text{NH}_4\text{HSO}_4 \div \text{KCl} = 2 \div 1$ increases, the general appearance of DTA curves and mass loss by TG changes (Fig. 2). The reaction mass is 304.82 g, and the calculated amount of released HCl is 36,5 g (11.974% of the charge mass).

The main phases of the cooled residue according to XRA are presented in Table 4.

The ammonium hydrosulfate increase in the charge leads to the formation of salts with more acidic properties, such as ammonium potassium hydrosulfate, in different stoichiometric ratios, depending on the excess of ammonium hydrosulfate and temperature.

According to the TG data (Fig. 2), at 350 °C, the formation of hydrogen chloride was only 73% of the calculated ones (Table 5).

Reaction 1 did not proceed to the end, considering the cooling stage, and the mass loss amounted to 9.93%. An additional amount of ammonia of 1.18% was released before the temperature reached 260 °C, since below that ammonia forms ammonium sulfate with ammonium hydrosulfate. A sufficient excess of ammonium hydrosulfate remains to form ammonium hydrosulfate and ammonium sulfate.

Table 4. Results of the X-ray phase analysis (XRA), cooled residue of reaction 1 in the temperature range from 350 °C to room temperature.

Phase Name	Formula	Figure of Merit	Phase Reg. Detail	DB Card Number
Dipotassium disulfate(VI)	$\text{K}_2(\text{S}_2\text{O}_7)$	1.335	ICDD (PDF-2 Release 2016 RDB)	01-077-4725
Potassium Ammonium Hydrogen Sulfate	$\text{K}(\text{NH}_4)\text{H}_2(\text{SO}_4)_2$	1.723	ICDD (PDF-2 Release 2016 RDB)	00-020-0850




Fig. (2). DTA and TG curves of reaction mixture 1 - ($\text{NH}_4\text{HSO}_4 + \text{KCl}$), at the molar ratio $\text{NH}_4\text{HSO}_4 \div \text{KCl} = 2 \div 1$.

Table 5. Results of mass loss by reactions by calculation and TG readings.

No.	Reaction Equations	$t_1 - t_2, ^\circ\text{C}$	$\Delta m_{\text{calculation}} \%$	$\Delta m_{\text{факт.}} \%$
0	$\text{NH}_4\text{HSO}_4(\text{s}) = \text{NH}_4\text{HSO}_4(\text{l})$	30 - 157,3	0	0
1	$\text{NH}_4\text{HSO}_4 + \text{KCl} = \text{KNH}_4\text{SO}_4 + \text{HCl}$	157 - 350	11.974	8.75
Cooling stage (XRA):				
15	$\text{NH}_4\text{HSO}_4 + \text{KNH}_4\text{SO}_4 = \text{NH}_4\text{K}(\text{HSO}_4)_2 + \text{NH}_3$	350 - 260 $^\circ\text{C}$		1,18
16	$\text{NH}_4\text{HSO}_4 + \text{NH}_3 = (\text{NH}_4)_2\text{SO}_4$	$\leq 260 ^\circ\text{C}$	-	
	Total:		11.974	9.93

Acid salts are likely to form in the presence of potassium sulfate in the reaction mixtures. The compounds of acid salts can be formed below the temperature of their thermal decomposition, both at the heating of reaction mixtures and at the cooling of condensed products of the direct reaction - metal sulfates and ammonium hydrosulfate (Table 3).

The reaction pyrolysis of ammonium sulfate is well-studied. The physicochemical methods show that the initial temperature point of the process is in the range of 235-260 $^\circ\text{C}$ [20, 26, 27], depending on the amount of suspension and heating rate at DTA. The mass loss values calculated from the reactions and TG curves comply efficiently (Table 3).

CONCLUSION

The following conclusions can be drawn from the results of the studies:

Thermodynamic functions of complex ammonium salts at 298K have been calculated using reference values of ionic increments.

The values of thermodynamic functions of supposed variants of chemical interaction of potassium salts with ammonium hydrosulfate melt have been calculated.

An analysis of the reaction of the interaction of potassium chloride with ammonium hydrosulfate was carried out, and it was found that:

- In the temperature range of 157 - 260 °C with equal molar ratios $\text{NH}_4\text{HSO}_4 \div \text{KCl}$, the condensed reaction product is a double salt - ammonium-potassium sulfate (NH_4KSO_4), and its intensive decomposition into constituent sulfates occurs above 260 °C with simultaneous pyrolysis of ammonium sulfate to complete formation of ammonium hydrosulfate and ammonia.

- The presence of ammonium hydrosulfate in the direct reaction products leads to the formation of potassium pyrosulfate. At the stage of cooling of condensed products of direct reaction in non-equilibrium conditions, depending on the stoichiometric ratios of components, the formation of hydrosulfates of different compositions is probable.

- In the incomplete pyrolysis of ammonium sulfate, the cooled residue contains $(\text{NH}_4)_3\text{HS}_2\text{O}_8$, i.e., the salt was formed from 2 ammonium sulfates: $(\text{NH}_4)_2\text{SO}_4 \cdot \text{NH}_4\text{HSO}_4$ and similarly, the product of incomplete decomposition of ammonium-potassium sulfate - NH_4KSO_4 is $\text{KHSO}_4 \cdot \text{K}_2\text{SO}_4$.

- The formation of hydrosulfates occurs in the presence of ammonium hydrosulfate in the composition of reaction mixtures, and the greater its proportion, the more acidic the double salts, the formation of which is accompanied by the simultaneous appearance of the main salt - ammonium sulfate;

- At a deficiency of ammonium hydrosulfate, on the contrary, the products of interaction are more basic salts.

- Increase of stoichiometric molar ratio of $\text{NH}_4\text{HSO}_4 \div \text{KCl}$ leads to the formation of more complex hydrated salts and ammonium sulfate.

Thus, the goal of obtaining ammonium-potassium sulfate fertilizer with three target components can be achieved in the form of ammonium potassium sulfate by holding the reaction mixture $\text{NH}_4\text{HSO}_4 + \text{KCl}$ below its decomposition temperature, preferably in the temperature range 150-200 °C.

AUTHOR'S CONTRIBUTIONS

All the authors contributed to the development of the concept and design of the study. The material's preparation, data collection and analysis were performed by Kalkaman Zhumashev, Akmaral Serikbayeva, Akkenzhe Busurmanova, Anar Akkenzheeva, Asia Boranbayeva, Zhansaule Altybayeva, Aigul Gusmanova, Aybala Narembekova, Feruza Berdikulova, Arman Mauleshev, and Tugelbai Kenbayev. The first version of the manuscript was written by Kalkaman Zhumashev and Akmaral Serikbayeva, and all authors commented on the previous versions of the manuscript. All authors read and approved the final manuscript. The authors declare that they have no conflict of interest.

LIST OF ABBREVIATIONS

TG	=	Thermogravimetry
DTA	=	Differential thermal analysis
DTG	=	Differential thermal gravimetry
cr.	=	Crystal

ΔH kCal	=	Enthalpy
ΔG	=	kCal - Gibbs energy
ΔS kCal	=	Entropy
T_{pb}	=	Temperature equilibrium
XRA	=	X-ray phase analysis
t_1-t_2 , °C	=	Temperature
$\Delta m_{\text{calculation}}$	=	Calculated weight difference
Δm_{fact}	=	Actual weight difference
s	=	Solid
l	=	Liquid

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

The data and supportive information are available within the article.

FUNDING

This research has been funded by the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan (Grant No. AP 19677917).

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] G. Schwenke, and G. McMullen, "Nitrogen volatilization from northern cropping soils", Available from: <https://grdc.com.au/resources-and-publications/grdc-update-papers/tab-content/grdc-update-papers/2009/09/nitrogen-volatilisation-from-northern-cropping-soils>
- [2] D. Alemneh, Z. Gete, A. Solomon, and L. Mekonnen, "Towards sustainable agriculture and rural development in the Ethiopian highlands", Available from: <https://www.fao.org/3/y5359e/y5359e00.htm>
- [3] S.L. Tisdale, "Sulphur in forage quality and ruminant nutrition", Available from: https://fertiliser-society.org/wp-content/uploads/2019/11/tsi_bulletin_22_ifs_reprint.pdf
- [4] R.G. Hoef, "Crop response to sulfur on Illinois soils", *J Fert Iss*, no. 2, pp. 94-105, 1985.
- [5] R. Dudal, "Forty years of soil fertility work in sub-Saharan Africa", In: *Integrated plant nutrient management in sub-Saharan Africa: From concept to practice*, CABI Publishing: Wallingford, UK, 2001, pp. 7-21. [<http://dx.doi.org/10.1079/9780851995762.0007>.]
- [6] H. Sen, M. Chien, and M. Gearhart, "Comparison of ammonium sulfate with other nitrogen and sulfur fertilizers in increasing crop production and minimizing environmental impact: A review", *Soil Sci.*, vol. 176, no. 7, 2011.
- [7] C.N. Kuznetsov, and S.V. Ardamakov, "Installation for the production of ammonium sulfate solution by direct mixing of sulfuric acid with ammonia", RU Patent 2393993
- [8] D.A. Pogaleshkin, A.S. Malyavin, A.M. Norov, and M.N. Tsykin, "A

- method for producing granular ammonium sulfate", RU Patent 2618763
- [9] D.A. Kosova, T.I. Navalayev, A.I. Maksimov, T.S. Babkina, and I.A. Uspenskaya, "Experimental investigation of the solid - Liquid phase equilibria in the water - Ammonium methanesulfonate and in the water - Sodium methanesulfonate systems", *Fluid Phase Equilib.*, vol. 443, pp. 23-31, 2017. [<http://dx.doi.org/10.1016/j.fluid.2017.04.006>]
- [10] M. Umarova, Z. Dekhkanov, and K. Aripov, "Obtaining potassium sulphate by conversion of potassium chloride and ammonium sulphate", *E3S Web Conf*, vol. 460, no. 4, 2023. [<http://dx.doi.org/10.1051/e3sconf/202346009035>]
- [11] P.K. Ghosh, P. M. Bhavnagar, and M. Ramniklal, US Patent 2015/0225250A1
- [12] D.A. Kosova, A.L. Emelina, M.A. Bykov, and A. Bykov, "Phase transitions of some sulfur-containing ammonium salts", *Thermochim. Acta*, vol. 595, pp. 61-66, 2014. [<http://dx.doi.org/10.1016/j.tca.2014.08.035>]
- [13] Y.H. Hotskyi, G.K. Ivanitsky, and A.R. Stepaniuk, "Investigation of the evaporation and crystallization of a sessile drop of ammonium sulfate solution on a smooth heated", *Thermophysics and Thermal Power Engineering*, vol. 43, no. 4, pp. 51-61, 2021. [<http://dx.doi.org/10.31472/ttpe.4.2021.6>]
- [14] A.R. Rao, P.S. Tide, B.K. George, and J. Mathew, "Quasi-dynamic model for dissolution coupled with reaction and precipitation of sodium bicarbonate in fed-batch reactive crystallization", *Chem. Eng. J. Adv.*, vol. 15, p. 100504, 2023. [<http://dx.doi.org/10.1016/j.cej.2023.100504>]
- [15] D.A. Makhrova, "Technology for the production of ammonium sulfate in chemical production", *XXIV International Scientific and practical Conference*, Anapa, Russia, 2021, pp. 26-30.
- [16] A-H.T. Kandil, M.F. Cheira, H.S. Gado, M.H. Soliman, H.M. Akl, H. Madiha, H. Soliman, and M.A. Hesham, "Ammonium sulfate preparation from phosphogypsum waste", *J. Rad. Res. Appl. Sci.*, vol. 10, no. 1, pp. 24-33, 2017. [<http://dx.doi.org/10.1016/j.jrras.2016.11.001>]
- [17] B.B. Sadykov, N.N. Volynskova, Sh.S. Namazov, and B.M. Beglov, "Technology for manufacturing fertilizer «superfos» containing nitrogen, phosphorus, sulfur and calcium", *Russ. J. Appl. Chem.*, vol. 83, no. 3, pp. 545-552, 2010.
- [18] V.V. Lapa, V.N. Emelyanova, and F.N. Leonov, *System fertilizer applications*, 2011. Available from: https://catalog.ggau.by/downloads/Publications/Jurgel/Jurgel_40.pdf
- [19] A. Marcu, D. Belis, and W. Stoefs, "Sectoral case study - Soda ash: Climate for sustainable growth",
- [20] K. Zhumashev, *Chemical and technological bases for improving ammonia-salt processes for the production of synthetic sodium carbonate.*, Monograph, 2023.
- [21] S.M. Kononov, A.N. Avramenko, O.A. Shirobokov, V.N. Gromotkov, and V.F. Perepechko, "Method for obtaining nitrogen-potassium fertilizer", RU Patent 2275347
- [22] A.I. Serebryakov, "Nitrogen-potassium fertilizer and method of its preparation", RU Patent 2198862, 2003.
- [23] A. Roine, "Outokumpu HSC chemistry for windows", Available from: www.outotec.com
- [24] B.P. Nikolsky, *Handbook of a chemist: Chemistry and chemical technology.*, vol. 21. 1966.
- [25] I.L. Knunyants, *Chemical encyclopedia*, vol. 1. Sovietskayaenciklopediya: Moscow, 1988, p. 623.
- [26] I.K. Thege, "DSC investigation of the thermal behaviour of (NH₄)₂SO₄, NH₄HSO₄ and NH₄NH₂SO₃", *Thermochim. Acta*, vol. 60, no. 2, pp. 149-159, 1983. [[http://dx.doi.org/10.1016/0040-6031\(83\)80265-2](http://dx.doi.org/10.1016/0040-6031(83)80265-2)]
- [27] N.N. Polyakov, N.M. Zhavoronkov, and E.Ya. Melnikov, "Ammonium sulfate, Handbook - M", *Chemistry*, pp. 228-234, 1987.

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.